

## University of Dayton eCommons

---

Mathematics Faculty Publications

Department of Mathematics

---

9-2010

# Coarser Connected Metrizable Topologies

Lynne Yengulalp

*University of Dayton*, [lyengulalp1@udayton.edu](mailto:lyengulalp1@udayton.edu)

Follow this and additional works at: [https://ecommons.udayton.edu/mth\\_fac\\_pub](https://ecommons.udayton.edu/mth_fac_pub)



Part of the [Applied Mathematics Commons](#), and the [Geometry and Topology Commons](#)

---

### eCommons Citation

Yengulalp, Lynne, "Coarser Connected Metrizable Topologies" (2010). *Mathematics Faculty Publications*. 40.  
[https://ecommons.udayton.edu/mth\\_fac\\_pub/40](https://ecommons.udayton.edu/mth_fac_pub/40)

This Article is brought to you for free and open access by the Department of Mathematics at eCommons. It has been accepted for inclusion in Mathematics Faculty Publications by an authorized administrator of eCommons. For more information, please contact [frice1@udayton.edu](mailto:frice1@udayton.edu), [mschlangen1@udayton.edu](mailto:mschlangen1@udayton.edu).

# Coarser connected metrizable topologies

Lynne Yengulalp<sup>1</sup>

*Department of Mathematics, University of Kansas, Lawrence, Kansas 66045, USA*

---

## Abstract

We show that every metric space,  $X$ , with  $w(X) \geq \mathfrak{c}$  has a coarser connected metrizable topology.

*Key words:* Coarser connected topology, extent, metrizable

*2008 MSC:* 54A10, 54A25, 54D05, 54D15, 54D70, 54E35

---

## 1. Introduction, Definitions, Notation

The study of coarser connected topologies was started by Tkačenko, Tkachuk, and Uspenskij who developed necessary and sufficient conditions for a topological space to have coarser connected Hausdorff or regular topology [5]. Continuing the development, Gruenhage, Tkachuk, and Wilson showed that any noncompact metrizable space has a coarser connected Hausdorff topology [4]. Fleissner, Porter and Roitman showed that any zero-dimensional metrizable space with weight at least  $\mathfrak{c}$  has a coarser connected metrizable topology [3]. We improve the last result by removing the zero-dimensional hypothesis. We show that every metric space with weight at least  $\mathfrak{c}$  has a coarser connected metrizable topology.

We write  $(X, \tau, \mu)$  for a metric space  $X$  with metric  $\mu$  and corresponding topology  $\tau$ . For a set  $A$  and cardinal  $\lambda$ , we write  $[A]^\lambda$  for the collection of subsets of  $A$  of cardinality  $\lambda$  and  $[A]^{<\lambda}$  for the collection on subsets of cardinality less than  $\lambda$ . The extent,  $e(X)$ , of a space  $X$  is the supremum of cardinals,  $\lambda$ , such that there is a closed discrete subset of  $X$  of size  $\lambda$ . If a space,  $X$ , has a closed discrete set of size  $e(X)$  we say the extent of  $X$  is attained in  $X$ .

Suppose  $U$  is a subset of a space  $X$ . If  $C$  is a closed discrete (with respect to  $U$ ) subset of  $U$ , then  $C$  is not necessarily closed in  $X$ . Hence, it is possible that  $e(U)$  is attained in  $U$ , but there is no closed discrete (in  $X$ ) subset of  $U$  of

---

<sup>1</sup>Department of Mathematics, University of Dayton, Dayton, Ohio 45469, USA  
yengullc@notes.udayton.edu

size  $e(U)$ . If, on the other hand, there is a closed discrete (in  $X$ ) subset of  $U$  of size  $e(U)$ , we say that  $e(U)$  is attained in  $X$ .

For a cardinal  $\kappa$  and positive real number  $\epsilon$ , the hedgehog with spininess  $\kappa$  and spine length  $\epsilon$  is the space  $Z = \{(\alpha, r) : \alpha \in \kappa, r \in (0, \epsilon]\} \cup \{(0, 0)\}$  with metric  $\rho$  defined as follows:

$$\rho((\alpha, r), (\alpha', r')) = \begin{cases} r + r' & \text{if } \alpha \neq \alpha' \\ |r - r'| & \text{if } \alpha = \alpha'. \end{cases}$$

The hedgehog space is a connected metric space.

## 2. Some Lemmas

If  $X$  is a metric space such that  $w(X) < \mathfrak{c}$ , then  $X$  does not necessarily have a coarser connected metrizable topology. For example the disjoint union of countably many Cantor sets is a separable, metrizable, disconnected space with no coarser connected regular topology [5]. We aim to show that all metric spaces,  $X$ , with  $w(X) \geq \mathfrak{c}$  have a coarser connected metrizable topology. This result has already been proven if  $e(X)$  is attained (Druzhinina [1], Fleissner, Porter, Roitman [3]). If  $e(X) = \mathfrak{c}$ , then we will see later that  $e(X)$  is attained. So, we consider metric spaces,  $X$ , such that  $e(X) > \mathfrak{c}$  is not attained.

In the proof of the main theorem, we partition the space  $(X, \tau, \mu)$  into subsets  $W_i$  for  $i \in \omega$  such that  $e(W_i)$  is attained in  $X$ . Lemma 2.1 will be used to define a coarser topology on  $X$  in which  $W_i$  is connected. We show that  $\mu$  can be adjusted so that  $W_i$  is connected without changing  $\mu$  on  $X \setminus W_i$ .

Lemmas 2.2–2.5 discuss the consequences of  $e(X)$  not being attained and provide a tool for the partition of  $X$  into  $W_i$ 's. We apply Lemma 2.6 to a particular compact subset of  $X$  for the purpose of defining the partition of  $X$  into  $W_i$ 's.

**Lemma 2.1.** *Suppose that  $Y$  is a subset of a metric space  $(X, \tau, \mu)$ ,  $e(\text{cl}Y) \geq \mathfrak{c}$  is attained in  $X$  by  $C \subset \text{int}(Y)$  and  $\text{diam}_\mu(\text{cl}Y) \leq \epsilon$ . If  $U \subset Y$  is an open set containing  $C$ , then there is a coarser topology  $\tau'$  on  $X$  generated by a corresponding metric  $\mu'$  such that*

- i)  $\mu' \leq \mu + 2\epsilon$ ,
- ii)  $\text{cl}_{\tau'}Y = \text{cl}_\tau Y$  and  $\tau'|_{\text{cl}Y}$  is connected, and
- iii)  $\mu|_{(X \setminus U)^2} = \mu'|_{(X \setminus U)^2}$ .

*Proof.* Let  $e(\text{cl}Y) = \kappa$ . So,  $C$  is closed discrete in  $X$  and  $|C| = \kappa$ . By replacing  $C$  with a subset, we may assume that  $|Y \setminus C| \geq \kappa$ . Let  $\mathcal{U} = \{U_c \subset U : c \in C\}$  be a discrete in  $X$  collection of open sets in  $X$  such that  $c \in$

$U_c \subset \text{cl}(U_c) \subset U$ . For each  $c \in C$  define a continuous function  $f_c : X \rightarrow [0, \epsilon]$  such that  $f_c(c) = \epsilon$  and  $f_c[X \setminus U_c] = \{0\}$ . For each  $x, y \in X$  define  $\mu^*(x, y) = \sum_{c \in C} |f_c(x) - f_c(y)| + \mu(x, y)$ . Because  $C$  is closed discrete in  $X$ , it is easy to check that  $\mu^*$  generates  $\tau$ ,  $\mu^*|_{(X \setminus U)^2} = \mu|_{(X \setminus U)^2}$  and  $\mu^* \leq \mu + 2\epsilon$ .

Let  $(Z, \rho)$  be a hedgehog space with spininess  $\kappa$  and spine length  $\epsilon$ . Let  $T = \{(\alpha, \epsilon) : \alpha \in \kappa\}$  and let  $S = Z \setminus T$ . The set  $T$  is the set of tips of the spines of the hedgehog which is a closed discrete set of size  $\kappa$ . Let  $D \subset Y \setminus C$  be a dense subset of  $Y \setminus C$  of size  $d(Y) = e(Y) = \kappa$ . Define a one-to-one map  $f : Z \rightarrow Y$  such that  $f[T] = D$  and  $f[S] = C$ . For  $x, y$  in the image of  $f$ , let  $\lambda(x, y) = \min\{\mu^*(x, y), \rho(f^{-1}(x), f^{-1}(y))\}$ . For all other  $x, y \in X$ , let  $\lambda(x, y) = \mu^*(x, y)$ .

Define a function  $\mu'$  on  $X \times X$  as follows:

$$\mu'(x, y) = \inf\{\lambda(x, x_1) + \lambda(x_1, x_2) + \cdots + \lambda(x_{n-1}, x_n) + \lambda(x_n, y)\}$$

where  $x_1, x_2, \dots, x_n$  ranges over all finite sequences (including the empty sequence) of distinct elements of  $X \setminus \{x, y\}$ . Since  $\rho$  and  $\mu^*$  satisfy the triangle inequality, in defining  $\mu'$  it suffices to consider sums of the form

$$\begin{aligned} \mu^*(x, x_1) + \rho(f^{-1}(x_1), f^{-1}(x_2)) + \cdots + \mu^*(x_{n-1}, x_n) \\ + \rho(f^{-1}(x_n), f^{-1}(y)) \end{aligned} \quad (1)$$

where the sum may start or end with either a  $\rho$  term or  $\mu^*$  term and the terms of the sum alternate between  $\rho$  and  $\mu^*$ . Since  $x_1, x_2, \dots, x_n$  are in the image of  $f$ , note that  $x_1, x_2, \dots, x_n \in D \cup C$ . Moreover, if  $x \notin Y$  and  $y \in Y$  each sum begins with a  $\mu^*$  term and  $x_1 \in Y$ . Hence,  $\mu'(x, y) \geq \mu^*(x, Y) \geq \mu(x, Y)$ . Letting  $y \in Y$  vary, we have  $\mu'(x, Y) \geq \mu(x, Y)$ .

Now we verify that  $\mu'$  is a metric. Since  $\mu'$  is an infimum over all finite sequences in  $X$ , it satisfies the triangle inequality. It is clear that  $\mu'(x, y) = \mu'(y, x)$  and  $\mu'(x, x) = 0$  for all  $x, y \in X$ . It remains to show that  $\mu'(x, y) = 0$  implies that  $x = y$ .

**Claim 1.** *Suppose that for a particular sum in the form of (1) and some  $m < n$ ,  $\lambda(x_i, x_{i+1}) < \epsilon$  for each  $i \in \{m, m+1, \dots, n-1\}$ . Then, if  $\lambda(x_m, x_{m+1}) = \rho(f^{-1}(x_m), f^{-1}(x_{m+1}))$  and  $x_m \in D$ , then  $x_{m+i} \in C$  for all  $i$  odd and  $x_{m+i} \in D$  for all  $i$  even in  $\{0, 1, \dots, n-m\}$ .*

*Proof.* Suppose that  $\lambda(x_m, x_{m+1}) = \rho(f^{-1}(x_m), f^{-1}(x_{m+1}))$  and  $x_m \in D$ . Also suppose that  $\lambda(x_i, x_{i+1}) < \epsilon$  for each  $i \in \{m, m+1, \dots, n-1\}$ . Fix  $i$  even such that  $0 \leq i < n-m$ , and suppose that  $x_{m+i} \in D$ . Since  $x_{m+i} \in D$ ,  $f^{-1}(x_{m+i}) \in T$ . But since the  $\rho$  and  $\mu$  terms of the sum alternate,

$$\lambda(x_{m+i}, x_{m+i+1}) = \rho(f^{-1}(x_{m+i}), f^{-1}(x_{m+i+1})) < \epsilon$$

and hence  $f^{-1}(x_{m+i+1}) \in S$ . By the definition of  $f$ ,  $x_{m+i+1} \in C$ . Now, since  $\lambda(x_{m+i+1}, x_{m+i+2}) = \mu^*(x_{m+i+1}, x_{m+i+2}) < \epsilon$  and  $x_{m+i+1} \in C$ , it must be that  $x_{m+i+2} \in D$ . So, since  $x_m \in D$ , by induction,  $x_{m+i} \in C$  for all  $i$  odd and  $x_{m+i} \in D$  for all  $i$  even in  $\{0, 1, \dots, n-m\}$ .  $\square$

**Claim 2.**  $\mu'(x, y) = 0$  implies that  $x = y$ .

*Proof.* Suppose there were  $x, y \in X$  such that  $\mu'(x, y) = 0$  but  $x \neq y$ . If defined, let  $x' = f^{-1}(x)$  and  $y' = f^{-1}(y)$ . Let  $x_1, \dots, x_n \in D \cup C$  be a sequence that yields an alternating  $\mu^*, \rho$  sum between  $x$  and  $y$  that is less than

$$\delta(x, y) = \min\{\epsilon, \mu^*(x, C \setminus \{x\}), \mu^*(y, C \setminus \{y\}), \rho(x', T \setminus \{x'\}), \rho(y', T \setminus \{y'\})\}.$$

Since  $\mu^*$  and  $\rho$  are metrics, we see that  $n \geq 1$ . Since  $C$  is closed discrete in  $X$  and  $T$  is closed in  $Z$ ,  $\delta(x, y)$  is a positive real number.

*Case 1.* The alternating sum begins and ends with  $\mu^*$  terms.

If  $x_1 \in C$  then  $\mu^*(x, x_1) \geq \mu^*(x, C \setminus \{x\}) \geq \delta(x, y)$ , which is a contradiction.

Note that  $n \geq 2$  is even. If  $x_1 \in D$ , then since  $\lambda(x_i, x_{i+1}) < \epsilon$  for each  $1 \leq i < n$  and  $\lambda(x_1, x_2) = \rho(f^{-1}(x_1), f^{-1}(x_2))$ , by Claim 1,  $x_n \in C$ . Hence  $\mu^*(x_n, y) \geq \mu^*(y, C \setminus \{y\}) \geq \delta(x, y)$ , which is a contradiction.

*Case 2.* The sum begins with a  $\mu^*$  term and ends with a  $\rho$  term.

If  $x_1 \in C$  then  $\mu^*(x, x_1) \geq \mu^*(x, C \setminus \{x\}) \geq \delta(x, y)$ , which is a contradiction.

Note that  $n$  is odd. If  $x_1 \in D$ , then since  $\lambda(x_i, x_{i+1}) < \epsilon$  for each  $1 \leq i < n$  and  $\lambda(x_1, x_2) = \rho(f^{-1}(x_1), f^{-1}(x_2))$ , by Claim 1,  $x_n \in D$ . Then  $\rho(f^{-1}(x_n), y') \geq \rho(y', T \setminus \{y'\}) \geq \delta(x, y)$ , which is a contradiction.

A symmetric argument shows that the sum beginning with  $\rho$  and ending with  $\mu^*$  yields a contradiction.

*Case 3.* The sum begins and ends with  $\rho$  terms.

Suppose  $x_1 \in D$ . Then  $\rho(f^{-1}(x), f^{-1}(x_1)) = \rho(x', f^{-1}(x_1)) \geq \rho(x', T \setminus \{x'\}) \geq \delta(x, y)$ , which is a contradiction.

Suppose  $x_1 \in C$ . Since  $\lambda(x_1, x_2) = \mu^*(x_1, x_2) < \epsilon$ ,  $x_2 \in D$ . Note that  $n \geq 2$  is even. If  $n = 2$ , then since  $x_2 \in D$ ,  $\rho(f^{-1}(x_n), y') \geq \rho(y', T \setminus \{y'\}) \geq \delta(x, y)$ , which is a contradiction. If  $n > 2$ ,  $\lambda(x_i, x_{i+1}) < \epsilon$  for each  $i$  with  $2 \leq i < n$  and  $\lambda(x_2, x_3) = \rho(f^{-1}(x_2), f^{-1}(x_3))$ . Therefore,  $x_n \in D$  by Claim 1. Hence  $\rho(f^{-1}(x_n), y') \geq \rho(y', T \setminus \{y'\}) \geq \delta(x, y)$ , which is a contradiction.  $\square$

So,  $\mu'$  defines a metric on  $X$ . Let  $\tau'$  be the topology on  $X$  generated by  $\mu'$ . Since  $\mu^*$  generates  $\tau$  and  $\mu' \leq \mu^*$ ,  $\tau'$  is coarser than  $\tau$ . Since  $\mu' \leq \mu^*$

and  $\mu^* \leq \mu + 2\epsilon$ , we have that  $\mu' \leq \mu + 2\epsilon$  showing i) of the lemma.

**Note 1.** In the proof of Claim 2, we have actually shown that for all  $x, y \in X$

$$\min\{\mu^*(x, y), \delta(x, y)\} \leq \mu'(x, y) \leq \mu^*(x, y).$$

**Claim 3.** *The map  $f : (Z, \rho) \rightarrow (Y, \mu'|_Y)$  is continuous.*

*Proof.* Let  $x \in Y$ ,  $\delta > 0$  and let  $U = B_{\mu'}(x, \delta)$  be the  $\mu'$ -ball of radius  $\delta$  about  $x$ . Suppose  $z \in f^{-1}[U]$ . Then  $\mu'(f(z), x) = \delta' < \delta$ . Let  $\xi = (\delta - \delta')/2$ . Suppose  $z' \in Z$  satisfies  $\rho(z, z') < \xi$ . We wish to show that  $\mu'(f(z'), x) < \delta$  so that  $B_\rho(z, \xi) \subset f^{-1}[U]$ . Since  $\mu'(x, f(z)) = \delta'$ , there is a sequence  $x_1, \dots, x_n \in C \cup D \subset Y$  such that

$$\lambda(x, x_1) + \dots + \lambda(x_n, f(z)) < \delta' + \xi.$$

Adding the term  $\rho(z, z') = \rho(f^{-1}(f(z)), f^{-1}(f(z')))$  to this sum,

$$\lambda(x, x_1) + \dots + \lambda(x_n, f(z)) + \rho(z, z') < \delta' + 2\xi < \delta$$

illustrates that the sequence  $x_1, \dots, x_n, f(z)$  yeilds a sum between  $x$  and  $f(z')$  that is less than  $\delta$ . Hence,  $\mu'(x, f(z')) < \delta$  as desired. So, the map  $f$  is continuous and the claim is proven.  $\square$

Since  $(Z, \rho)$  is connected,  $\tau \subset \tau'$  and  $C \cup D$  is dense in  $(Y, \tau|_Y)$ , by Claim 3,  $(Y, \mu'|_Y)$  is connected. For  $x \notin Y$ , we argued before that  $\mu'(x, Y) \geq \mu(x, Y)$ , so  $\text{cl}_{\tau'} Y \subset \text{cl}_\tau Y$ . However  $\text{cl}_\tau Y \subset \text{cl}_{\tau'} Y$ , since  $\tau'$  is coarser than  $\tau$ . Hence  $\text{cl}_{\tau'} Y = \text{cl}_\tau Y = \text{cl}_\tau(D \cup C)$  and therefore  $(\text{cl} Y, \tau'|_{\text{cl} Y})$  is connected. Therefore, ii) of the lemma is shown.

We now show iii). Let  $x, y \notin U$ . We verify that  $\mu'(x, y) = \mu(x, y)$ . By definition,  $\mu'(x, y) \leq \mu^*(x, y) = \mu(x, y)$  so we need only show that  $\mu'(x, y) \geq \mu^*(x, y)$ .

Suppose for a contradiction that  $\mu'(x, y) < \mu^*(x, y)$ . In other words, there exist  $x_1, \dots, x_n \in D \cup C$  such that

$$\mu'(x, y) \leq \lambda(x, x_1) + \dots + \lambda(x_n, y) < \mu^*(x, y).$$

Since  $x, y \notin U$ ,  $\mu^*(x, y) = \mu(x, y)$ . If  $f^{-1}(x)$  and  $f^{-1}(y)$  are defined, then since  $x, y \notin C$ ,  $\rho(f^{-1}(x), f^{-1}(y)) = 2\epsilon > \mu(x, y) = \mu^*(x, y)$ . Hence, the sequence  $x_1, \dots, x_n$  is not empty. Now, since  $\text{diam}_\mu(Y) \leq \epsilon$ , for any  $z_1, z_2 \in Y$ ,

$$\mu(x, y) \leq \mu(x, z_1) + \epsilon + \mu(z_2, y).$$

Hence,

$$\mu(x, y) \leq \mu(x, Y) + \epsilon + \mu(y, Y).$$

Combining these inequalities we have

$$\mu'(x, y) \leq \lambda(x, x_1) + \cdots + \lambda(x_n, y) < \mu(x, Y) + \epsilon + \mu(y, Y). \quad (2)$$

*Case 1.*  $x, y \in Y$ .

Then inequality (2) becomes  $\mu'(x, y) \leq \lambda(x, x_1) + \cdots + \lambda(x_n, y) < \epsilon$ . Since  $x, y \notin U$ ,  $\mu^*(x, C \setminus \{x\}) = \mu^*(x, C) \geq \epsilon$  and  $\mu^*(y, C \setminus \{y\}) = \mu^*(y, C) \geq \epsilon$ . Since  $x \notin C$ , if  $x' = f^{-1}(x)$  is defined, then  $x \in D$ . Hence  $\rho(x', T \setminus \{x'\}) \geq \epsilon$ . Similarly, if  $y' = f^{-1}(y)$  is defined,  $\rho(y', T \setminus \{y'\}) \geq \epsilon$ . So,  $\delta(x, y) = \min\{\epsilon, \mu^*(x, C \setminus \{x\}), \mu^*(y, C \setminus \{y\}), \rho(x', T \setminus \{x'\}), \rho(y', T \setminus \{y'\})\} \geq \epsilon$ . By Note 1 after Claim 2,  $\mu'(x, y) \geq \min\{\mu^*(x, y), \epsilon\}$ . This contradicts the assumption that  $\mu'(x, y) < \mu^*(x, y)$ .

*Case 2.*  $x \notin Y$  and  $y \in Y$ .

Since  $y \in Y$ ,  $\mu(y, Y) = 0$ , hence (2) becomes  $\mu'(x, y) \leq \lambda(x, x_1) + \lambda(x_1, x_2) + \cdots + \lambda(x_n, y) < \mu(x, Y) + \epsilon$ . Suppose  $x_1 \in C$ . Since  $x \notin Y$ ,  $\lambda(x, x_1) = \mu^*(x, x_1) = \mu(x, x_1) + \epsilon \geq \mu(x, Y) + \epsilon$ , a contradiction. Suppose that  $x_1 \in D$ . Since  $\lambda(x, x_1) = \mu^*(x, x_1) \geq \mu(x, Y)$  we have that  $\lambda(x_1, x_2) + \cdots + \lambda(x_n, y) < \epsilon$ . If  $n$  is even,  $\lambda(x_n, y) = \mu^*(x_n, y)$  and by Claim 1,  $x_n \in C$ . Therefore  $\lambda(x_n, y) = \mu^*(x_n, y) = \mu(x_n, y) + \epsilon > \epsilon$ , a contradiction. If  $n$  is odd,  $\lambda(x_n, y) = \rho(f^{-1}(x_n), f^{-1}(y))$  and so  $y \in C \cup D$ . Since  $y \notin U$ , we have that  $y \in D$  and therefore  $\rho(f^{-1}(x_n), f^{-1}(y)) = 2\epsilon > \epsilon$ , a contradiction.

A symmetric argument yields a contradiction in the case that  $x \in Y$  and  $y \notin Y$ .

*Case 3.*  $x, y \notin Y$ .

Since  $x, y \notin Y$ , they are not in the image of  $f$  and therefore the alternating sum must start and end with  $\mu^*$  terms implying  $n$  is even and  $n \geq 2$ . Since  $x_1, x_n \in C \cup D \subset Y$ ,  $\mu^*(x, x_1) \geq \mu(x, x_1) \geq \mu(x, Y)$  and  $\mu^*(x_n, y) \geq \mu(x_n, y) \geq \mu(y, Y)$ . Combining this with (2) gives:

$$\begin{aligned} \rho(f^{-1}(x_1), f^{-1}(x_2)) + \mu^*(x_2, x_3) + \cdots + \rho(f^{-1}(x_{n-1}), f^{-1}(x_n)) &< \epsilon, \\ \mu^*(x, x_1) &< \mu(x, Y) + \epsilon \end{aligned} \quad (3)$$

and

$$\mu^*(x_n, y) < \mu(y, Y) + \epsilon. \quad (4)$$

Suppose  $x_1 \in C$ . Then  $\mu^*(x, x_1) = \mu(x, x_1) + \epsilon \geq \mu(x, Y) + \epsilon$  contradicting (3). So,  $x_1 \notin C$ . Similarly,  $x_n \in C$  contradicts (4). So,  $x_n \notin C$ . Hence we have shown that  $x_1, x_n \in D$ . Now,  $\lambda(x_i, x_{i+1}) < \epsilon$  for each  $i$  with  $1 \leq i < n$  and  $\lambda(x_1, x_2) = \rho(f^{-1}(x_1), f^{-1}(x_2))$ . So, since  $x_1 \in D$  and  $n$  is even, by Claim 1,  $x_n \in C$ , which is a contradiction. Therefore,  $\mu'(x, y) = \mu^*(x, y) = \mu(x, y)$ , and iii) of the lemma follows.  $\square$

If  $X$  is a metric space with  $e(X)$  not attained, then  $X$  must have a certain form. This form is described in the next lemma.

**Lemma 2.2.** (Fitzpatrick, Gruenhage, Ott [2]) *Let  $(X, \tau)$  be a metric space with metric  $\mu$  in which  $e(X) = \kappa$  is not attained. Let  $K$  be the set of points  $x$  of  $X$  such that every neighborhood of  $x$  has extent  $\kappa$ . Then*

- (1)  $\kappa$  is a singular cardinal of cofinality  $\omega$ .
- (2)  $K$  is a compact, nowhere dense subset of  $X$ .
- (3) If  $U$  is an open subset of  $X$  such that  $\text{cl}_\tau U \cap K = \emptyset$ , then  $e(U) < \kappa$ .
- (4)  $K$  is nonempty.

Recall that König's Lemma implies that  $\text{cf}(\mathfrak{c}) > \omega$ . So, if  $(X, \tau)$  is a metric space and  $e(X) = \mathfrak{c}$ , by Lemma 2.2, the extent of  $X$  must be attained.

**Definition 2.3.** (see [3]) An open set  $V$  is called *e-homogeneous* if for every nonempty open subset  $V'$  of  $V$ ,  $e(V') = e(V)$ .

Suppose  $U$  is a nonempty open subset of a metric space. We argue that there is a nonempty open e-homogeneous set  $V$  such that  $V \subset U$ . If  $U$  is e-homogeneous then set  $V = U$ . If  $U$  is not e-homogeneous, let  $V_0 = U$  and let  $V_1$  be a nonempty open subset of  $U$  such that  $e(V_1) < e(U)$ . Continue by defining  $V_n \subset V_{n-1}$  such that  $e(V_n) < e(V_{n-1})$  if  $V_{n-1}$  is not homogeneous. Since  $e(V_1), e(V_2), \dots$  is a decreasing sequence of cardinal numbers, there is  $n \in \omega$  such that  $V = V_n$  is e-homogeneous.

Suppose the extent of an open subset,  $U$ , of a metric space is not attained in  $U$ . Apply Lemma 2.2 with  $X = U$ . Then, by property (2), there is  $x \in U \setminus K$  and an open set  $V \subset U$  with  $e(V) < e(U)$  and  $x \in V$ . Therefore, if  $U$  is an e-homogeneous subset of a metric space,  $e(U)$  is attained in  $U$ . However, as mentioned before, this does not mean that  $e(U)$  is attained in  $X$ .

**Lemma 2.4.** *Let  $U$  be an e-homogeneous subset of a metric space  $(X, \tau, \mu)$  with metric  $\mu$  and  $e(U) > \aleph_0$ . Then  $e(U)$  is attained in  $X$ .*

*Proof.* Since  $U$  is e-homogeneous,  $e(U) = \lambda$  is attained in  $U$ . Suppose that  $\text{cf}(\lambda) > \aleph_0$ . Then,  $e(U)$  is attained by some closed discrete (in  $U$ ) subset  $C' \subset U$  of cardinality  $\lambda$ . However, it may not be the case that  $C'$  is closed discrete in  $X$ . We will construct a set  $C \subset U$  such that  $C$  is closed discrete in  $\text{cl}_\tau U$ , hence closed discrete in  $X$ , and  $|C| = e(U) = \lambda$ . Let  $\mathcal{U}$  be a discrete collection of open subsets of  $U$  that witnesses that  $C'$  is closed discrete in  $U$ . Let

$$L = \text{cl}_\tau \left( \bigcup \mathcal{U} \right) \setminus \bigcup_{U \in \mathcal{U}} \text{cl}_\tau U.$$

Note that  $L \subset \text{cl}_\tau U \setminus U$ . If  $L = \emptyset$  then  $\mathcal{U}$  is discrete in  $X$  and  $C'$  is closed discrete in  $X$ . Therefore, suppose that  $L \neq \emptyset$ . Let  $C_n = \{c \in C' : \mu(c, L) \geq$



$1/n\}$  and note that  $\bigcup C_n = C'$ . Since  $cf(\lambda) > \aleph_0$  there is  $n \in \omega$  so that  $|C_n| = \lambda$ . Set  $C = C_n$ . By construction,  $C$  is closed and  $\text{cl}_\tau C \cap L = \emptyset$ , so  $C$  is closed discrete in  $X$ ,  $C \subset U$  and  $|C| = \lambda = e(U)$  as desired.

Now suppose that  $cf(\lambda) = \aleph_0$ . Let  $\{\lambda_n : n \in \omega\}$  be an increasing sequence of cardinals such that  $\lambda = \sup_{n \in \omega} \lambda_n$ . Let  $W$  be an open subset of  $U$  such that  $\text{cl}_\tau W \subset U$ . Note that since  $U$  is e-homogeneous,  $e(W) = e(\text{cl}_\tau W) = e(U) = \lambda$ . Since  $e(\text{cl}_\tau W) = \lambda > \aleph_0$  there is  $A \subset \text{cl}_\tau W \subset U$  a countable closed discrete set in  $\text{cl}_\tau W$ , hence closed discrete in  $X$ . Let  $\mathcal{U} = \{U_n : n \in \omega\}$  be a discrete collection of open subsets of  $U$  that witnesses  $A$  is discrete in  $X$ . For each  $n \in \omega$ ,  $e(\text{cl}_\tau U_n) = \lambda > \lambda_n$ , so there is  $C_n \subset \text{cl}_\tau U_n$  a closed discrete in  $X$  subset of cardinality  $\lambda_n$ . Let  $C = \bigcup C_n$ . Since  $C_n$  is closed discrete in  $\text{cl}_\tau U_n$ , it is closed discrete in  $X$ . Moreover, since  $\mathcal{U}$  is discrete,  $C$  is closed discrete in  $X$ . Finally,  $|C| = \lambda = e(\text{cl}_\tau U)$  by construction and since  $U_n \subset U$  for each  $n$ ,  $C \subset U$ . In either case,  $|C| = \lambda = e(U)$ ,  $C \subset U$  and  $C$  is closed and discrete in  $X$ . Hence,  $e(U)$  is attained in  $X$ .  $\square$

**Lemma 2.5.** *Let  $(X, \tau)$  be a metric space with metric  $\mu$  in which  $e(X) = \kappa$  is not attained. Let  $K$  be the set of points  $x$  of  $X$  such that every neighborhood of  $x$  has extent  $\kappa$ . Then, for every open set  $U$  meeting  $K$  and every  $\theta < \kappa$  there is an open subset  $V$  of  $U$  such that  $\text{cl}_\tau V \subset U$ ,  $e(V) > \theta$  is attained in  $X$  and  $K \cap \text{cl}_\tau V = \emptyset$ .*

*Proof.* Let  $\mathcal{V}$  be a maximal pairwise disjoint collection of e-homogeneous subsets  $V$  of  $U \setminus K$  such that  $\text{cl}_\tau V \cap K = \emptyset$ . For each  $V \in \mathcal{V}$ , since  $\text{cl}_\tau V \cap K = \emptyset$ , by item (3) of Lemma 2.2,  $e(V) < \kappa$ . Suppose that for some  $V \in \mathcal{V}$ ,  $e(V) > \theta$ . By Lemma 2.4,  $e(V)$  is attained in  $X$  and we are done. Suppose on the other hand that  $e(V') \leq \theta$  for all  $V' \in \mathcal{V}$ . Since  $\mathcal{V}$  is maximal,  $K$  is nowhere dense and every open subset of  $X$  has an e-homogeneous subset,  $\bigcup \mathcal{V}$  is dense in  $U$ . Since  $X$  is metric,  $e(W) = d(W)$  for any open subset  $W$ . Suppose that  $|\mathcal{V}| = \lambda < \kappa$ . Then,  $d(U) = d(\bigcup \mathcal{V}) \leq \lambda \cdot \theta < \kappa$  which is a contradiction. So,  $|\mathcal{V}| = \kappa$ . Since  $\mu(V', K) > 0$  for all  $V' \in \mathcal{V}$  and  $cf(\kappa) = \omega$ , there is  $n \in \omega$  such that  $|\{V' \in \mathcal{V} : \mu(K, V') > 1/n\}| > \theta$ . Set  $\mathcal{V}' = \{V' \in \mathcal{V} : \mu(K, V') > 1/n\}$ . We now refine  $\mathcal{V}'$  to a discrete collection of size  $> \theta$ . Let

$$L = \text{cl}_\tau(\bigcup \mathcal{V}') \setminus \bigcup_{V' \in \mathcal{V}'} \text{cl}_\tau V'$$

and let  $L_\epsilon = \{x \in U : \mu(x, L) \leq \epsilon\}$ . Since  $\bigcap_{\epsilon > 0} L_\epsilon = L$  and  $V' \cap L = \emptyset$  for

all  $V' \in \mathcal{V}'$ , there is  $m \in \omega$  such that  $|\{V' \in \mathcal{V}' : V' \setminus L_{1/m} \neq \emptyset\}| > \theta$ . Set  $\mathcal{V}'' = \{V' \in \mathcal{V}' : V' \setminus L_{1/m} \neq \emptyset\}$ . For each  $V' \in \mathcal{V}''$ , let  $W(V') = V' \setminus L_{1/m}$ . The collection  $\mathcal{W} = \{W(V') : V' \in \mathcal{V}''\}$  by construction is discrete and has

cardinality  $> \theta$ . Set  $V = \bigcup \mathcal{W}$ . For each  $W \in \mathcal{W}$ , choose  $x_W \in W$ . Let  $C = \{x_W : W \in \mathcal{W}\}$ . Then  $C$  is closed discrete in  $X$  since  $\mathcal{W}$  is discrete in  $X$ , and  $|C| = |\mathcal{W}| > \theta$ . Since  $e(V) \leq \sup\{e(W) : W \in \mathcal{W}\} \cdot |\mathcal{W}| \leq \theta \cdot |\mathcal{W}| = |C|$ , we have that  $e(V) > \theta$  is attained in  $X$ .  $\square$

**Lemma 2.6.** *Let  $K$  be a compact metric space with topology  $\sigma$  and let  $\mathcal{U} \in [\sigma]^{<\omega}$  be a pairwise disjoint collection such that  $\bigcup \mathcal{U}$  is dense in  $K$ . If  $\epsilon > 0$  and  $\mathcal{V}$  is the collection of open subsets of  $K$  with diameter less than  $\epsilon$ , then there exists a pairwise disjoint  $\mathcal{V}' \in [\mathcal{V}]^{<\omega}$  such that  $\mathcal{V}'$  refines  $\mathcal{U}$  and  $\bigcup \mathcal{V}'$  is dense in  $K$ .*

*Proof.* Since  $\mathcal{V}$  covers  $K$ , compact, there are  $n \in \omega$  and  $V_1, V_2, \dots, V_n \in \mathcal{V}$  such that  $K = \bigcup \{V_i : 1 \leq i \leq n\}$ . Define  $\hat{V}_1 = V_1$  and for  $1 < i \leq n$  let  $\hat{V}_i = V_i \setminus \text{cl}(\bigcup \{V_j : 1 \leq j < i\})$ . Let  $\hat{\mathcal{V}} = \{\hat{V}_i : 1 \leq i \leq n\}$ . Note that  $\hat{\mathcal{V}}$  is pairwise disjoint and  $\bigcup \hat{\mathcal{V}}$  is dense in  $X$ . Now define  $\mathcal{V}' = \{V \cap U : V \in \hat{\mathcal{V}}, U \in \mathcal{U}\}$ . Since  $V \cap U \subset U$  for each  $V \in \hat{\mathcal{V}}$  and  $U \in \mathcal{U}$ ,  $\mathcal{V}'$  refines  $\mathcal{U}$ . Since  $\text{diam}(V \cap U) \leq \text{diam}(V) \leq V_i$  for some  $1 \leq i \leq n$ ,  $\text{diam}(V') < \epsilon$  for all  $V' \in \mathcal{V}'$ . Hence  $\mathcal{V}' \subset \mathcal{V}$ . Since  $\hat{\mathcal{V}}$  is pairwise disjoint,  $\mathcal{V}'$  is as well. Finally,  $\bigcup \mathcal{V}' = \bigcup \hat{\mathcal{V}} \cap \bigcup \mathcal{U}$  and therefore  $\bigcup \mathcal{V}'$  is dense in  $K$ , since  $\bigcup \mathcal{V}'$  and  $\bigcup \mathcal{U}$  are open and dense in  $K$ .  $\square$

### 3. The Main Theorem

**Theorem 3.1.** *If  $(X, \tau)$  is a metric space with metric  $\mu$  and  $w(X) = \kappa \geq \mathfrak{c}$ , then there is  $\sigma$ , a topology on  $X$  coarser than  $\tau$ , such that  $(X, \sigma)$  is connected and metrizable.*

*Proof.* As mentioned in the beginning of Section 2, the theorem has been proven in the case when  $e(X)$  is attained. Therefore, we assume  $e(X) > \mathfrak{c}$  is not attained. Re-scale  $\mu$  so that  $\text{diam}_\mu(X) < 1/2$  by replacing it with  $\frac{\mu}{2(1+\mu)}$ . Let  $K$  be the set of points  $x$  of  $X$  such that every neighborhood of  $x$  has extent  $\kappa$ . By Lemma 2.2,  $K$  is compact.

Let  $\mathcal{C}_0^* = \{K\}$ . For each  $n \in \omega \setminus \{0\}$ , define  $\mathcal{C}_n^* \subset \tau|_K$ , a pairwise disjoint finite collection with the following properties.

- $\text{cl}(\bigcup \mathcal{C}_n^*) = K$ .
- $\mathcal{C}_n^*$  refines  $\mathcal{C}_{n-1}^*$ .
- $B \in \mathcal{C}_n^*$  implies  $\text{diam}(B) < 1/2^n$ .

Let  $n \in \omega \setminus \{0\}$ . Apply Lemma 2.6 with  $\sigma = \tau|_K$ ,  $\mathcal{U} = \mathcal{C}_{n-1}^*$  and  $\epsilon = 1/2^n$  to get  $\mathcal{V}'$ , a pairwise disjoint collection of open sets in  $K$  with diameter less than  $1/2^n$  that refines  $\mathcal{C}_{n-1}^*$  and whose union is dense in  $K$ . Set  $\mathcal{C}_n^* = \mathcal{V}'$ .

### Definition of $B_i$ 's

For  $n \in \omega$  enumerate the elements of  $\mathcal{C}_n^*$  as  $\mathcal{C}_n^* = \{B_i^* : i_n \leq i < i_{n+1}\}$  with an increasing sequence of integers,  $i_n$ , where  $i_0 = 0$ . For each  $i \in \omega$ , let  $L_i = \{x \in X : \mu(x, B_i^*) \leq \mu(x, K \setminus B_i^*)\}$ , the set of points in  $X$  that are closer to  $B_i^*$  than they are to its complement. Fix  $n \in \omega$  and let  $B_{i_n} = L_{i_n}$  and for each  $i$  with  $i_n < i < i_{n+1}$ , let  $B_i = L_i \setminus \bigcup \{\text{cl}(B_j) : i_n \leq j < i\}$ . Note that  $B_0^* = K$ ,  $B_0 = X$  and that  $B_j^* \subset B_i^*$  implies  $L_j \subset L_i$ .

We define  $\mathcal{C}_n = \{B_i : i_n \leq i < i_{n+1}\}$  and verify the following.

- (i) For each  $n \in \omega$ ,  $\mathcal{C}_n$  is pairwise disjoint.
- (ii) For each  $n \in \omega$ ,  $\bigcup \mathcal{C}_n$  is dense in  $X$ .
- (iii) For all  $i \in \omega$ ,  $\text{int}(B_i) \cap K \neq \emptyset$ .

From the definition of  $B_i$  and  $\mathcal{C}_n$ , (i) is clear. Towards (ii), let  $n \in \omega$  and let  $x \in X$ . Since  $\bigcup \mathcal{C}_n^*$  is dense in  $K$ ,  $\mu(x, K) = \mu(x, \bigcup \mathcal{C}_n^*)$ . Since  $\mathcal{C}_n^*$  is finite, there is  $i$  such that  $i_n \leq i < i_{n+1}$  and  $\mu(x, K) = \mu(x, B_i^*)$ . Therefore,  $\mu(x, B_i^*) \leq \mu(x, K \setminus B_i^*)$  and either  $x \in B_i$  or  $x \in \text{cl}(\bigcup \{B_j : i_n \leq j < i\})$ . In either case,  $x \in \text{cl}(\bigcup \mathcal{C}_n)$ . For (iii), note that  $\text{int}(B_i) \cap K = B_i^* \setminus \text{cl}(\bigcup \{B_j^* : i_n \leq j < i\}) = B_i^* \neq \emptyset$  since  $\mathcal{C}_n^*$  is pairwise disjoint.

Let  $U_i = \{x \in X : \mu(x, K) < 1/2^{i+1}\}$ .

**Claim 1.** For  $n \in \omega$ ,  $i_n \leq i < i_{n+1}$ ,  $\text{diam}_\mu(L_i \cap U_m) \leq 3/2^n$  for any  $m \geq n - 1$ .

*Proof.* Note that if  $x \in L_i$ ,  $\mu(x, B_i^*) \leq \mu(x, K \setminus B_i^*)$  which implies  $\mu(x, K) = \mu(x, B_i^*)$ . For  $i$  with  $i_n \leq i < i_{n+1}$ ,  $m \geq n - 1$ ,  $x, y \in L_i \cap U_m$  and  $\epsilon > 0$  there exist  $x_0, y_0 \in B_i^*$  such that  $\mu(x, x_0), \mu(y, y_0) < 1/2^{m+1} + \epsilon/2 \leq 1/2^n + \epsilon/2$ . So,

$$\mu(x, y) \leq \mu(x, x_0) + \mu(y, y_0) + \mu(x_0, y_0) \leq 2/2^n + \epsilon + 1/2^n \leq 3/2^n + \epsilon.$$

Hence,  $\text{diam}_\mu(L_i \cap U_m) \leq 3/2^n$ .  $\square$

### Definition of $W_i$ 's

We define  $\mathcal{W} = \{W_i : i \in \omega\}$ , a pairwise disjoint collection of open subsets of  $X$  such that

- (i)  $\text{cl}(W_i) \cap K = \emptyset$ ,
- (ii)  $e(W_i) > \mathfrak{c}$  is attained in  $X$  by  $C_i \subsetneq W_i$ ,
- (iii) for  $i_n \leq i < i_{n+1}$ ,  $\text{diam}_\mu(W_i) \leq 3/2^n$  and
- (iv)  $\bigcup \mathcal{W}$  is dense in  $X$ .

Let  $\hat{W}_0 = \emptyset$ . Apply Lemma 2.5 with  $\theta = \mathfrak{c}$  and  $U = \text{int}(B_0) = X$  to get an open subset  $V$  of  $X$  such that  $e(V) > \theta = \mathfrak{c}$  is attained in  $X$  by  $C_0 \subset V$  and  $\text{cl}_\tau V \cap K = \emptyset$ . By replacing  $C_0$  with a subset of the same cardinality, we may assume that  $C_0 \subsetneq V$ . Set  $W_0 = V$  and  $S_0 = V$ . Let  $k_0 = \min\{k \geq 1 : S_0 \subset X \setminus U_k\} \geq 1$ . By definition  $W_0$  is open, and since  $\text{cl}V \cap K = \text{cl}(W_0) \cap K = \emptyset$ , (i) holds. By the choice of  $W_0$ , (ii) holds and (iii) is trivial since  $\text{diam}_\mu(X) \leq 1/2$ .

Suppose we have defined  $W_i$  for all  $i$  with  $0 \leq i < i_n$  so that (i), (ii), and (iii) are satisfied. Also suppose that  $S_m = \bigcup\{W_i : i < i_{m+1}\}$  is dense in  $X \setminus \text{cl}(U_{k_{m-1}+1})$  and that  $k_m = \min\{k \geq m : S_m \subset X \setminus U_k\}$  whenever  $0 < m < n$ .

Let  $i$  be such that  $i_n \leq i < i_{n+1}$ . Let  $\hat{W}_i = B_i \setminus (\text{cl}(U_{k_{n-1}+1} \cup S_{n-1}))$ . Since  $\text{int}(B_i) \cap K \neq \emptyset$  and  $K \subset X \setminus \text{cl}(S_{n-1})$ ,  $\text{int}(B_i) \setminus \text{cl}(S_{n-1})$  is an open set meeting  $K$ . So, apply Lemma 2.5 with  $\theta = \max\{e(\hat{W}_i), \mathfrak{c}\}$  and  $U = \text{int}(B_i) \setminus \text{cl}(S_{n-1})$  to get an open subset  $V$  of  $U$  such that  $\text{cl}_\tau V \subset U$ ,  $e(V) > \theta$  is attained in  $X$  by  $C_i \subset V$  and  $\text{cl}_\tau V \cap K = \emptyset$ . As before, we may assume  $C_i \subsetneq V$ . Set  $W_i = \hat{W}_i \cup V$ . According to our choice,  $W_i$  satisfies (i) and (ii) for each  $i$  such that  $i_n \leq i < i_{n+1}$ . Set  $S_n = \bigcup\{W_i : i_n \leq i < i_{n+1}\} \cup S_{n-1}$ . Let  $k_n = \min\{k \geq n : S_n \subset X \setminus U_k\}$ . By Claim 1,  $\text{diam}(B_i \cap U_{n-1}) \leq 3/2^n$  since  $B_i \subset L_i$  and  $i_n \leq i < i_{n+1}$ . Also,  $S_{n-1}$  is dense in  $X \setminus U_{k_{n-2}+1}$ . Therefore, for any  $i$  as above,  $W_i = \hat{W}_i \cup V \subset B_i \setminus \text{cl}(S_{n-1}) \subset B_i \cap U_{k_{n-2}+1} \subset B_i \cap U_{n-1}$ , since  $k_{n-2} \geq n-2$ . Hence  $\text{diam}_\mu(W_i) \leq 3/2^n$  and (iii) is satisfied.

Towards (iv), since  $K$  is nowhere dense in  $X$ , we only show that  $\bigcup \mathcal{W}$  is dense in  $X \setminus K$ . Let  $x \in X \setminus K$  and let  $n \in \omega \setminus \{0\}$  be such that  $x \in X \setminus \text{cl}(U_{k_{n-1}+1})$ . Then either  $x \in \text{cl}(S_{n-1}) \subset \text{cl}(S_n)$  or  $x \in X \setminus \text{cl}(U_{k_{n-1}+1} \cup S_{n-1})$ . If  $x \in X \setminus \text{cl}(U_{k_{n-1}+1} \cup S_{n-1})$  then since  $\{B_i : i_n \leq i < i_{n+1}\}$  is dense in  $X$ ,  $x \in \text{cl}(B_i \setminus (\text{cl}(U_{k_{n-1}+1} \cup S_{n-1}))) \subset \text{cl}(W_i) \subset \text{cl}(S_n)$  for some  $i$ . In either case  $x \in \text{cl}(S_n)$ . So,

$$x \in \text{cl}\left(\bigcup\{W_i : i \in \omega\}\right) = \text{cl}\left(\bigcup\{S_n : n \in \omega\}\right).$$

Hence  $\bigcup\{W_i : i \in \omega\}$  is dense in  $X \setminus K$  and in  $X$ .

#### Linking the $W_i$ 's

Suppose that  $n, i, j \in \omega$  are such that  $i_n \leq i < i_{n+1} \leq j < i_{n+2}$  and  $B_j^* \subset B_i^*$ . Then  $W_i \subset B_i \cap U_{n-1} \subset L_i \cap U_{n-1}$  and  $W_j \subset B_j \cap U_n \subset L_j \cap U_n \subset L_i \cap U_{n-1}$ . Hence by Claim 1,  $x \in W_i$  and  $y \in W_j$  implies  $\mu(x, y) \leq 3/2^n$ . Since for each  $j \in \omega$ ,  $C_j \subsetneq W_j$ , we may choose  $x_j \in W_j \setminus C_j$  arbitrarily. For every  $i$  with  $i_n \leq i < i_{n+1}$ , let  $J_i = \{j : i_{n+1} \leq j < i_{n+2}, B_j^* \subset B_i^*\}$  and let  $X_i = \{x_j : j \in J_i\}$ . Notice that  $\text{diam}(W_i \cup X_i) \leq 3/2^n$ ,  $C_i \subset W_i \subset \text{int}(W_i \cup X_i)$  and that by the definition of  $B^*$ ,  $\bigcup_{i \in \omega} J_i = \omega \setminus \{0\}$ .

### Defining the connected topology on $X$

We define a sequence of metrics  $\mu_n$  on  $X$  such that  $\nu = \lim \mu_n$  is a well defined metric that generates a coarser connected topology on  $X$ . We define  $\mu_n$  by induction.

#### Stage 0

Apply Lemma 2.1 with  $Y = W_0 \cup X_0$ , metric  $\mu$ , topology  $\tau$ , closed discrete set  $C_0$ ,  $\epsilon = 3/2$  and  $U = W_0$  to get a topology  $\tau_0$  on  $X$  generated by the metric  $\mu_0$  on  $X$  such that

- (i)  $\mu_0 \leq \mu + 6/2$ ,
- (ii)  $\text{cl}_\tau(W_0 \cup X_0) = \text{cl}_{\tau_0}(W_0 \cup X_0)$  and  $\tau_0|_{W_0 \cup X_0}$  is connected, and
- (iii)  $\mu|_{(X \setminus W_0)^2} = \mu_0|_{(X \setminus W_0)^2}$ .

Note that since  $X_0$  is finite, property (ii) implies that  $\text{cl}_{\tau_0}(W_0) = \text{cl}_{\tau_0}(W_0 \cup X_0)$ . So, for all  $i \neq 0$ ,  $\text{int}_{\tau_0}(W_i) = W_i \setminus X_0 \supset W_i \setminus \{x_i\}$  and therefore since  $x_i \notin C_i$ ,  $C_i \subset \text{int}_{\tau_0}(W_i)$ .

#### Stage $n \geq 1$

Set  $\sigma_0 = \tau_{n-1}$  and  $\rho_0 = \mu_{n-1}$ . For each  $i$  such that  $i_n \leq i < i_{n+1}$ , let  $j = i - i_n$  and apply Lemma 2.1 with  $Y = W_i \cup X_i$ ,  $C = C_i$ ,  $\tau = \sigma_j$ ,  $\mu = \rho_j$ ,  $U = \text{int}_{\sigma_j}(W_i)$  and  $\epsilon = 3/2^n$  to get a topology  $\sigma_{j+1}$  on  $X$  generated by the metric  $\rho_{j+1}$  on  $X$  such that

- (i)  $\rho_{j+1} \leq \rho_j + 6/2^n$ ,
- (ii)  $\text{cl}_{\sigma_j}(W_i \cup X_i) = \text{cl}_{\sigma_{j+1}}(W_i \cup X_i)$  and  $\sigma_{j+1}|_{W_i \cup X_i}$  is connected, and
- (iii)  $\rho_j|_{(X \setminus \text{int}_{\sigma_j}(W_i))^2} = \rho_{j+1}|_{(X \setminus \text{int}_{\sigma_j}(W_i))^2}$ .

Let  $\mu_n = \rho_{m_n}$  and  $\tau_n = \sigma_{m_n}$ . Note that, as in Stage 0, property (ii) implies  $C_i \subset \text{int}_{\tau_n}(W_i)$  for all  $i \geq i_{n+1}$ . It is a consequence of (i) and (iii), that if  $x \in X \setminus \bigcup\{W_i : i \leq i_{n+1}\}$  and  $y \in \bigcup\{W_i : i < i_{n+1}\}$  then  $\mu_n(x, y) \leq \mu(x, y) + 6/2^n$ . Furthermore,  $\mu_n = \mu_{n-1}$  on  $X \setminus \bigcup\{W_i : i_n \leq i < i_{n+1}\}$  and  $\tau_n|_{W_i \cup X_i}$  is connected for each  $i$  such that  $i_n \leq i < i_{n+1}$ .

Define  $\nu(x, y) = \lim \mu_n(x, y)$ . This map is a well defined metric since for any  $x, y \in X$ ,  $\nu(x, y) = \mu_m(x, y)$  for all

$$m \geq \max\{n : i_n \leq i < i_{n+1} \text{ and } x \in W_i \text{ or } y \in W_i\} \cup \{0\}.$$

Let  $\sigma$  be the topology generated by  $\nu$ . To show that  $\sigma \subset \tau$  we show that  $\nu$  preserves convergent  $\mu$  sequences. Suppose that  $\{y_j : j \in \omega\}$  and  $y$  are

such that  $\lim_{j \rightarrow \infty} \mu(y, y_j) = 0$ . If  $y \notin K$  there are  $n, m \in \omega$  such that  $y, y_j \in X \setminus \text{cl}(U_{k_n+1})$  for all  $j \geq m$ . Then,  $\nu(y, y_j) = \mu_n(y, y_j)$  for all  $j \geq m$ , hence  $\lim_{j \rightarrow \infty} \nu(y, y_j) = \lim_{j \rightarrow \infty} \mu_n(y, y_j) = 0$ , since  $\mu_n$  preserves  $\mu$  convergent sequences. Now suppose that  $y \in K$ . If  $y_j \in U_m$  for all  $m$ , let  $m_j = j$ . Otherwise, let  $m_j = \max\{m : y_j \in U_m\}$ . Since  $\mu(y, y_j) \rightarrow 0$ ,  $m_j \rightarrow \infty$ . If there is  $i \in \omega$  such that  $y_j \in W_i$  then there is  $n$  such that  $i \geq i_n \geq n = m_j$ , since  $y_j \in W_i \subset U_{m_j}$ . In this case, by the consequence of (i) and (iii),  $\nu(y, y_j) \leq \mu(y, y_j) + 6/2^n = \mu(y, y_j) + 6/2^{m_j}$ . If  $y_j \notin W_i$  for all  $i \in \omega$  then  $\nu(y, y_j) = \mu(y, y_j)$ . In either case  $\nu(y, y_j) \leq \mu(y, y_j) + 6/2^{m_j}$  and therefore  $\nu(y, y_j) \rightarrow 0$ . Note that  $\nu|_{(W_i \cup X_i)^2} = \mu_n|_{(W_i \cup X_i)^2}$  for  $i_n \leq i < i_{n+1}$ . Hence  $W_i \cup X_i$  is connected in  $\sigma$ . Furthermore,  $X_i \cap W_j \neq \emptyset$  for each  $j \in J_i$  and hence  $W_i \cup \bigcup_{j \in J_i} W_j$  is connected as well. This means that  $W_i$  is ‘linked’ to  $W_0$  for every  $i$  such that  $1 = i_1 \leq i < i_2$  and since  $\bigcup_{i \in \omega} J_i = \omega \setminus \{0\}$ , any later  $W_i$  is ‘linked’ to  $W_0$ . Therefore any  $\sigma$ -clopen subset,  $Z$ , of  $X$  would have to be empty, or contain  $W_i$  for all  $i$ . Since  $\bigcup_{i \in \omega} W_i$  is dense in  $X$ ,  $Z$  is trivial. Hence  $\sigma$  is connected.  $\square$

This work is part of my Ph.D. dissertation, University of Kansas, 2009.

#### 4. Acknowledgements

I would like to thank the referee for the many helpful suggestions. They have greatly improved the paper. I would also like to thank my advisor, William Fleissner, for his guidance and support.

#### References

- [1] I. Druzhinina, Condensations onto connected metrizable spaces, *Houston Journal of Mathematics* 30 (3) (2004) 751–766.
- [2] B. Fitzpatrick, G. Gruenhage, J. Ott, Topological completions of metrizable spaces, *Proceedings of the American Mathematical Society*, Vol. 117, No. 1 (Jan 1993) 259–267.
- [3] W. Fleissner, J. Porter, J. Roitman, Coarser connected topologies, *Topology Appl.* 142 (2004) 131–157.
- [4] G. Gruenhage, V.V. Tkachuk, R.G. Wilson, Weaker connected and weaker nowhere locally compact topologies for metrizable and similar spaces, *Topology Appl.* 120 (2002) 365–384.
- [5] M.G. Tkačenko, V.V. Tkachuk, V.V. Uspenskij, R.G. Wilson, In quest of weaker connected topologies, *Comment. Math. Univ. Carolin.* 37(4) (1996) 825–841.
- [6] L. Yengulalp, Coarser connected topologies and non-normality points, (2009) Ph.D. dissertation, University of Kansas.